

SOME ANALYTICAL RESULTS ON CLASSICAL TUNNELING OF SOMMERFELD PARTICLE

Alexander A. Vlasov

High Energy and Quantum Theory
Department of Physics
Moscow State University
Moscow, 119899
Russia

A simple example of especially constructed potential barrier enables to show analytically (not numerically) the existence of tunneling effect for a Sommerfeld particle.

03.50.De

Let us remind that Sommerfeld model of charged rigid sphere [1] (Sommerfeld particle) is the simplest theoretical model to take into consideration the "back-reaction" of self-electromagnetic field on equation of motion of a radiating extended charged body (for point-like charge we have the known Lorentz-Dirac equation with all its problems: renormalization of mass, preacceleration, runaway solutions, etc.).

For the case of simplicity here we consider the nonrelativistic, linear in velocity, version of Sommerfeld model [2].

So let the total charge of a uniformly charged sphere be Q , mechanical mass - m , radius - a . Then its equation of motion reads [1,2]:

$$m\vec{v} = \vec{F}_{ext} + \eta [\vec{v}(t - 2a/c) - \vec{v}(t)] \quad (1)$$

here $\eta = \frac{Q^2}{3ca^2}$, $\vec{v} = d\vec{R}/dt$, $\vec{R}$ - coordinate of the center of the shell, $\vec{F}_{ext}$ - some external force.

This equation has no runaway solutions and solutions with preacceleration. But, as was shown in [3], it has the solutions, which one can interpret as classical tunneling. The physics of such effect is simple - Sommerfeld particle due to retardation begins to "feel" the action of potential barrier too late, when this barrier is overcome.

In [3] the existence of solutions with tunneling effect was demonstrated with the help of numerical calculations.

Here we present one simple problem, which enables us to see the appearance of classical tunneling without numerical calculations but using only analytical considerations.

Let the Sommerfeld particle move along z -axis ($\vec{R} = (0, 0, R)$) in a static

electric field $\vec{E} = (0, 0, E_z)$ produced by electric potential ϕ in a form of a step:

$$\phi = B[\theta(z) - \theta(z - S)], \quad E_z = -\frac{d\phi}{dz} = -B\delta(z) + B\delta(z - S) \quad (2)$$

Then the external force F_{ext} , acting on rigid sphere with density of charge ρ ,

$$\rho = \frac{Q}{4\pi a^2} \delta(|\vec{r} - \vec{R}| - a)$$

reads

$$F_{ext} = \int d\vec{r} \rho E_z = \frac{QB}{4\pi a^2} \int d\vec{r} \delta(|\vec{r} - \vec{R}| - a) [-\delta(z) + \delta(z - S)]$$

With new variables $\vec{r} \equiv \vec{\xi} + \vec{R}$, $d\vec{r} = d\vec{\xi} = \xi^2 d\xi \sin \theta d\theta d\phi$, substituting $\cos \theta \equiv \mu$ (it leads to $z = R + \xi\mu$), integrating over ϕ and ξ , we get

$$F_{ext} = \frac{QB}{2a} \int_{-1}^{+1} d\mu [-\delta(R + a\mu) + \delta(R + a\mu - S)] = \frac{QB}{2a} \left[- \int_{R-a}^{R+a} dw \delta(w) + \int_{R-a-S}^{R+a-S} dw \delta(w) \right]$$

If $S \geq a$ then the expression in square brackets yields the result

0 for $R \leq -a$, -1 for $-a \leq R \leq +a$, 0 for $+a \leq R \leq S-a$, +1 for $S-a \leq R \leq S+a$, 0 for $R \geq S+a$

After this, eq.(1) in dimensionless variables $y = R/L$, $\tau = ct/L$, $\delta = 2a/L$, for the relations $S = L = 2a$ (taken for simplicity), is reduced to

$$\frac{d^2 y}{d\tau^2} = k \left[\frac{dy(\tau - 1)}{d\tau} - \frac{dy(\tau)}{d\tau} \right] + \lambda \Phi \quad (3)$$

here $k = \frac{2Q^2}{3mc^2 a}$, $\lambda = \frac{QB}{mc^2}$,
and

$$\Phi = \begin{cases} 0, & y < -1/2, \\ -1, & -1/2 < y < 1/2, \\ +1, & 1/2 < y < 3/2, \\ 0, & 3/2 < y, \end{cases}$$

Let us note that in the limit of zero particle's size $a \rightarrow 0$ the force F_{ext} tends to two delta-functions:

$$\lim_{a \rightarrow 0} F_{ext} = QB[-\delta(R) + \delta(R - S)]$$

This expression gives the following Newtonian eq. of motion for point-like particle

$$m\dot{v} = QB[-\delta(R) + \delta(R - S)] = -QB \frac{d}{dR} [-\theta(R) + \theta(R - S)] \quad (4)$$

Equation (4) has the first integral:

$$m \frac{v^2}{2} + QB[\theta(R) - \theta(R - S)] = \text{const} = m \frac{v_0^2}{2}$$

Thus for $0 < R < S$ we have

$$v = \sqrt{v_0^2 - (2QB/m)}$$

This solution describes the overcoming of a point-like particle the potential barrier only if the initial velocity v_0 is greater then the critical value v_{cr} , equal to $2QB/m$ or in dimensionless form

$$\frac{v_{cr}}{c} \equiv \dot{y}_{cr} = \sqrt{2\lambda} \quad (5)$$

If $k \equiv 0$ then the equation (3) for $-1/2 < y < +1/2$, $0 < \tau < t_1$ has the solution

$$y = -1/2 + v_0\tau - \lambda\tau^2/2, \quad \dot{y} = v_0 - \lambda\tau \quad (6)$$

It follows from (6) that the point $y = +1/2$ is achieved (if the velocity $\dot{y}$ is always positive) at the moment of time t_1 :

$$t_1 = \frac{v_0 - \sqrt{(v_0)^2 - 2\lambda}}{\lambda} \quad (7)$$

Consequently the potential barrier is overcome if $v_0 > \sqrt{2\lambda}$, and the minimal initial velocity for this is $v_0 = \sqrt{2\lambda}$ with $\dot{y}(t_1) = 0$ and

$$t_1 = v_0/\lambda = 2/v_0 \quad (8)$$

(the positive part of the force in the equation (3) for $1/2 < y < 3/2$ does not hinder the motion of the particle, so for analysis of conditions for overcoming the barrier one can consider only the interval: $-1/2 < y < +1/2$),

so for $t_1 = 2$, one must take $v_0 = 1$ and $\lambda = 1/2$, and for $t_1 = 3$ — $v_0 = 2/3$ and $\lambda = 2/9$ and so on.

If k in (3) is not equal to zero and if the point $y = 1/2$ is achieved at the moment of time $\tau = t_1$ in such a way that $\dot{y}(t_1) = 0$ (this means that in the preceding moments of time the velocity is positive) then the further motion of particle, $1/2 < y < 3/2$, is governed by the equation

$$\frac{d^2y}{d\tau^2} + k \frac{dy(\tau)}{d\tau} = k \frac{dy(\tau - 1)}{d\tau} + \lambda \equiv f$$

where $f > 0$.

This eq. with the above condition $\dot{y}(t_1) = 0$ has the solution

$$\dot{y} = e^{(-k\tau)} \int_{t_1}^{\tau} dt' f(t') e^{(kt')} > 0$$

consequently the barrier is overcome and the particle continues its motion for $\tau > t_1$.

Taking all this into consideration, let's construct the " tunneling" solutions of the equation (3).

Let the Sommerfeld particle reach the point $y = -1/2$ at the moment of time $\tau = 0$, $y = 1/2$ — at $\tau = t_1$.

Let the particle's velocity for $\tau < 0$ be constant:

$$\frac{dy}{d\tau} = v_0 = \text{const}$$

Let the condition be true: $\dot{y}(t_1) = 0$ (i.e. the barrier is overcome) and let

$$t_1 = 2 \quad (9)$$

Then one must integrate the eq. (3) into two stages: in the first, for the interval $0 < \tau < 1$, $-1/2 < y < y_1$ and, in the second, for the interval $1 < \tau < 2$, $y_1 < y < 2$.

Solution on the first interval is

$$\dot{y} = v_0 - \frac{\lambda}{k} + \frac{\lambda}{k} e^{(-k\tau)} \quad (10)$$

and on the second —

$$\begin{aligned} \dot{y} &= v_0 - 2\frac{\lambda}{k} + a_0 e^{(-k\tau)} + \frac{\lambda}{k} \tau e^{(-k\tau)}, \\ a_0 &= \frac{\lambda}{k} (e^{(k\tau)} + 1) - \lambda e^{(k\tau)} \end{aligned} \quad (11)$$

Conditions $y(t_1 = 2) = 1/2$ $\dot{y}(t_1 = 2) = 0$ are reduced to the set

$$\begin{aligned} v_0 &= f_1 \lambda \\ \lambda &= 1/f_2 \end{aligned}$$

here

$$f_1 = \frac{2}{k} - e^{(-k)} \left(1 + \frac{1}{k} + \frac{e^{(-k)}}{k}\right) \quad (12)$$

$$f_2 = \frac{1 - e^{(-k)}}{k} + \frac{3 - 2e^{(-k)} - e^{(-2k)}}{k^2} - 2e^{(-k)} \left(1 + \frac{1}{k} + \frac{e^{(-k)}}{k}\right) \quad (13)$$

The analysis of the functions $f_1(k)$, $f_2(k)$ (12, 13) shows that the condition $v_0 < \sqrt{2\lambda}$ for the solution (10, 11) is valid due to the inequality $f_1/\sqrt{2f_2} < 1$ - i.e. there is the tunneling - the initial velocity has the forbidden, from newtonian point of view, values, but ought to the inequality $f_1/f_2 > 1$ all of them are greater than 1: $v_0 > 1$ (for $k \equiv 0$, in accordance with (8), $v_0 = 1$ and $\lambda = 1/2$).

Let now instead of (9) be

$$t_1 = 3 \quad (14)$$

Then one must integrate the eq. (3) into tree stages: in the first, for the interval $0 < \tau < 1$, $-1/2 < y < y_1$ with the solution in the form (10), in the second, for the interval $1 < \tau < 2$, $y_1 < y < y_2$ with the solution in the form (11), and in the third, for the interval $2 < \tau < 3$, $y_2 < y < 1/2$ with the solution in the form

$$\begin{aligned} \dot{y} &= v_0 - 3\frac{\lambda}{k} + e^{(-kt)}(a_1 + ke^{(k)}(a_0 - \lambda e^{(k)})\tau + k\lambda e^{(2k)}\tau^2/2), \\ a_1 &= e^{(2k)}\left(\frac{\lambda}{k} - 2\lambda + 2k\lambda - \lambda e^{(-k)} + \frac{\lambda}{k}e^{(-k)}\right) \end{aligned} \quad (15)$$

Conditions $y(t_1 = 3) = 1/2$ and $\dot{y}(t_1 = 3) = 0$ are reduced to the set

$$v_0 = f_3\lambda$$

$$\lambda = 1/f_4$$

here

$$f_3 = \frac{e^{(-3k)}}{2k}(6e^{(3k)} - (k^2 + 2k + 2)e^{(2k)} - (4k + 2)e^{(k)} - 2) \quad (16)$$

$$f_4 = \frac{e^{(-3k)}}{2k^2}((6k+12)e^{(3k)} - (3k^3+7k^2+10k+6)e^{(2k)} - (12k^2+10k+4)e^{(k)} - 6k-2) \quad (17)$$

Analysis of functions (16, 17) shows that the condition $v_0 < \sqrt{2\lambda}$ for the solution (15) is valid — $f_3/\sqrt{2f_4} < 1$ - i.e. there is the tunneling, and, contrary to the above case, the range of values of v_0 is: $2/3 < v_0 < 1$ (for $k \equiv 0$ in accordance with (8), $v_0 = 2/3$ and $\lambda = 2/9$).

The construction of tunneling solutions one can continue for $t_1 = 4, 5, 6, \dots$ and so on. It is obviously that the greater is the value of t_1 , the lower is the value of the minimal velocity v_0 necessary to overcome the barrier, but the more is complicated the form of the solution.

REFERENCES

1. A.Sommerfeld, Gottingen Nachrichten, 29 (1904), 363 (1904), 201 (1905).
2. L.Page, Phys.Rev., 11, 377 (1918)
T.Erber, Fortschr. Phys., 9, 343 (1961)
P.Pearle in "Electromagnetism",ed. D.Tepliz, (Plenum, N.Y., 1982), p.211.
A.Yaghjian, "Relativistic Dynamics of a Charged Sphere". Lecture Notes in Physics, 11 (Springer-Verlag, Berlin, 1992).
F.Rohrlich, Am.J.Phys., 65(11), 1051 (1997). Phys.Rev., D60, 084017 (1999).
3. Alexander A. Vlasov, physics/9911059.